

## WHAT IS CLAIMED IS:

1. A substantially pure or isolated polypeptide comprising a segment exhibiting sequence homology to a corresponding portion of a mature protein selected from the group consisting of:
- i) TECK;
  - ii) MIP-3 $\alpha$ ;
  - iii) MIP-3 $\beta$ ;
  - iv) DC CR; and
  - v) M/DC CR;
- wherein said homology is at least about 70% identity and said portion is at least about 25 amino acids.
2. The protein of Claim 1, further comprising a second segment exhibiting:
- a) at least about 90% identity over at least 9 amino acids; or
  - b) at least about 80% identity over at least 17 amino acids.
3. The polypeptide of Claim 1, wherein said polypeptide:
- a) is from a warm blooded animal selected from the group of birds and mammals, including a mouse or human;
  - b) comprises a natural sequence from Tables 1 through 5;
  - c) exhibits a post-translational modification pattern distinct from a natural form of said polypeptide;
  - d) is made by expression of a recombinant nucleic acid;
  - e) comprises synthetic sequence;
  - f) is detectably labeled;
  - g) is conjugated to a solid substrate;
  - h) is conjugated to another chemical moiety;
  - i) is a fusion protein;

- 5 j) is in a denatured conformation, including  
detergent denaturation;  
k) further comprises an epitope tag;  
l) is an immunogenic polypeptide;  
m) has a defined homogeneous molecular weight;  
n) is useful as a carbon source;  
o) is an allelic variant of SEQ ID NO: 2, 4, 6, 8,  
10, or 12;  
p) is a 3-fold or less substituted form of a natural  
10 sequence;  
q) is in a sterile composition;  
r) is in a buffered solution or suspension;  
s) is in a regulated release device;  
t) comprises a post-translational modification;  
15 u) is in a cell; or  
v) is in a kit which further comprises instructions  
for use or disposal of reagents therein.

4. An isolated or recombinant nucleic acid encoding  
20 said protein of Claim 1, where said portion consists of  
sequence from the coding region of SEQ ID NO: 1, 3, 5, 7,  
9, or 11.

5. The nucleic acid of Claim 4, wherein said nucleic  
25 acid:

- a) exhibits at least about 80% identity to a natural  
cDNA encoding said segment;  
b) is in an expression vector;  
c) further comprises a promoter;  
30 d) further comprises an origin of replication;  
e) is from a natural source;  
f) is detectably labeled;  
g) comprises synthetic nucleotide sequence;  
h) is less than 6 kb;  
35 i) is from a mammal;  
j) comprises a natural full length mature coding  
sequence;

- 5      k) is in a kit, which also comprises instructions for  
         use or disposal of reagents therein;  
         l) is a specific hybridization probe for a gene  
         encoding said protein;  
         m) is a PCR product; or  
         n) is in a cell.

10      6.      A method of using a purified nucleic acid of  
         Claim 5, comprising a step of expressing said nucleic acid  
         to produce a protein.

15      7.      An isolated or recombinant nucleic acid which  
         encodes at least eight consecutive residues of SEQ ID NO:  
         2, 4, 6, 8, 10, or 12.

20      8.      The nucleic acid of Claim 7, which encodes at  
         least:

- 25      a) twelve consecutive residues from SEQ ID NO: 2, and  
         further comprises a coding sequence of at least  
         17 nucleotides from SEQ ID NO: 1;  
         b) twelve consecutive residues from SEQ ID NO: 4, and  
         further comprises a coding sequence of at least  
         17 nucleotides from SEQ ID NO: 3;  
         c) twelve consecutive residues from SEQ ID NO: 6, and  
         further comprises a coding sequence of at least  
         17 nucleotides from SEQ ID NO: 5;  
         d) twelve consecutive residues from SEQ ID NO: 8, and  
         further comprises a coding sequence of at least  
         17 nucleotides from SEQ ID NO: 7;  
30      e) twelve consecutive residues from SEQ ID NO: 10,  
         and further comprises a coding sequence of at  
         least 17 nucleotides from SEQ ID NO: 9; or  
         f) twelve consecutive residues from SEQ ID NO: 12,  
         and further comprises a coding sequence of at  
35      least 17 nucleotides from SEQ ID NO: 11.

9. The nucleic acid of Claim 7, wherein said nucleic acid:

- a) exhibits at least about 80% identity to a natural cDNA encoding said segment;
- 5 b) is in an expression vector;
- c) further comprises a promoter;
- d) further comprises an origin of replication;
- e) encodes a 3-fold or less substituted sequence from a natural sequence;
- 10 f) is from a natural source;
- g) is detectably labeled;
- h) comprises synthetic nucleotide sequence;
- i) is less than 6 kb;
- j) is from a mammal;
- 15 k) is attached to a solid substrate, including in a Southern or Northern blot;
- l) comprises a natural full length coding sequence;
- m) is in a cell; or
- 20 n) is in a detection kit, which also comprises instructions for use or disposal of reagents therein.

10. A nucleic acid which hybridizes under stringent wash conditions of 55° C and less than 150 mM salt to the  
25 nucleic acid of Claim 7.

11. The nucleic acid of Claim 10, which exhibits at least about 85% identity over a stretch of at least about 30 nucleotides to a primate sequence of SEQ ID NO: 1, 3, 5,  
30 7, 9, or 11.

12. The nucleic acid of Claim 10, wherein:  
a) said identity is at least 90%; or  
b) said stretch is at least 75 nucleotides.

35

13. The nucleic acid of Claim 10, wherein:  
a) said identity is at least 95%; or

b) said stretch is at least 100 nucleotides.

14. A binding compound comprising an antigen binding fragment from an antibody which binds to a protein of Claim

5 1.

15. The binding compound of Claim 14, wherein:

- a) said polypeptide is a mouse or human protein;
- b) said antibody is raised against a mature peptide  
10 sequence of Tables 1 through 5;
- c) said antibody is a monoclonal antibody;
- d) said binding compound is attached to a solid  
substrate;
- e) said binding compound is in a sterile composition;
- 15 f) said binding compound binds to a denatured  
antigen, including a detergent denatured antigen;
- g) said binding compound is detectably labeled;
- h) said binding compound is an Fv, Fab, or Fab2  
fragment;
- 20 i) said binding compound is conjugated to a chemical  
moiety;
- j) said binding compound is in a detection kit which  
also comprises instructions for use or disposal  
of reagents therein.

25

16. A cell which makes said antibody of Claim 14.

17. A method of purifying a polypeptide using a binding compound of Claim 14 to specifically separate said  
30 polypeptides from others.

18. A method of generating an antigen-binding compound complex comprising the step of contacting a sample comprising said antigen to a sample comprising a binding  
35 compound of Claim 14.

19. A method of modulating physiology or development of a cell expressing a receptor for a chemokine selected from the group selected from:

- 5           a) TECK;  
          b) MIP-3 $\alpha$ ; or  
          c) MIP-3 $\beta$ ;

comprising contacting said cell with a composition comprising:

- 10           i) an agonist or mutein of said chemokine; or  
          ii) an antibody antagonist of said chemokine.

20. The method of Claim 19, wherein said cell is a macrophage or lymphocyte.

15 21. The method of Claim 19, wherein said physiology is selected from:

- a) a cellular calcium flux;  
          b) a chemoattractant response;  
          c) cellular morphology modification responses;  
20          d) phosphoinositide lipid turnover; or  
          e) an antiviral response.

22. The method of Claim 19, wherein:

- 25          a) said receptor is DC CR and said chemokine is MIP-3 $\alpha$ ;  
          b) said physiology is pulmonary physiology; or  
          c) said cell is an eosinophil.